

# Notes on the Stochastic Exponential and Logarithm\*

Martin Larsson<sup>†</sup>

Johannes Ruf<sup>‡</sup>

February 20, 2017

## Abstract

Stochastic exponentials are defined for semimartingales on stochastic intervals, and stochastic logarithms are defined for nonnegative semimartingales, up to the first time the semimartingale hits zero continuously. In the case of (nonnegative) local supermartingales, these two stochastic transformations are inverse to each other. The reciprocal of a stochastic exponential is again a stochastic exponential on a stochastic interval.

**Keywords:** Stochastic exponential, stochastic logarithm, stochastic interval, involution.

**MSC2010 subject classification:** Primary 60G99; secondary: 60H10, 60H99.

## 1 Introduction

This note collects results for the calculus of stochastic exponentials and logarithms of semimartingales, possibly defined on stochastic intervals only. We have found these results useful in a number of situations, and while they are no doubt well known, we were not able to find a suitable reference. Stochastic exponentials for not necessarily continuous semimartingales date back to at least Doléans-Dade (1976). Besides basically any textbook on stochastic calculus, the survey article Rheinländer (2010) reviews well known properties of stochastic exponentials of semimartingales (defined on the whole positive axis). Kallsen and Shiryaev (2002) provides further interesting identities, especially relating to exponential and logarithmic transforms, a subject which we do not discuss in this note.

We proceed as follows. In Section 2, we establish notation and introduce the concept of processes on stochastic intervals. In Section 3, we define stochastic exponentials and logarithms, discuss their basic properties, and prove that they are inverse to each other. In

---

\*Parts of this note appeared in the unpublished manuscript Larsson and Ruf (2014). J.R. acknowledges generous support from the Oxford-Man Institute of Quantitative Finance, University of Oxford.

<sup>†</sup>Department of Mathematics, ETH Zurich, Rämistrasse 101, CH-8092, Zurich, Switzerland. E-mail: martin.larsson@math.ethz.ch

<sup>‡</sup>Department of Mathematics, London School of Economics and Political Science, Columbia House, Houghton St, London WC2A 2AE, United Kingdom. E-mail: j.ruf@lse.ac.uk

Section 4, we describe the stochastic logarithm of the reciprocal of a stochastic exponential. Finally, in Section 5, we provide some examples. These examples illustrate that stochastic exponentials of semimartingales, defined on stochastic intervals only, arise naturally.

## 2 Notation and processes on stochastic intervals

The following definitions are consistent with those in Jacod and Shiryaev (2003), to which the reader is referred for further details. We work on a stochastic basis  $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ , where the filtration  $\mathbb{F} = (\mathcal{F}_t)_{t \geq 0}$  is right-continuous but not necessarily augmented with the  $\mathbb{P}$ -nullsets. Relations between random quantities are understood in the almost sure sense.

Given a process  $X = (X_t)_{t \geq 0}$ , write  $X_-$  for the left limit process (limit inferior if a limit does not exist) and  $\Delta X = X - X_-$  for its jump process, using the convention  $X_{0-} = X_0$ . The corresponding jump measure is denoted by  $\mu^X$ , and for any (random) function  $F : \Omega \times \mathbb{R}_+ \times \mathbb{R} \rightarrow \mathbb{R}$ , the stochastic integral of  $F$  with respect to  $\mu^X$  is the process  $F * \mu^X$  given by

$$F * \mu_t^X = \begin{cases} \sum_{s \leq t} F(s, \Delta X_s) \mathbf{1}_{\{\Delta X_s \neq 0\}}, & \text{if } \sum_{s \leq t} |F(s, \Delta X_s)| \mathbf{1}_{\{\Delta X_s \neq 0\}} < \infty, \\ +\infty, & \text{otherwise,} \end{cases} \quad t \geq 0.$$

Semimartingales are required by definition to be right-continuous, almost surely admitting left limits. If  $X$  is a semimartingale,  $X^c$  denotes its continuous local martingale part, and  $H \cdot X$  is the stochastic integral of an  $X$ -integrable process  $H$  with respect to  $X$ .

For a stopping time  $\tau$ , we let  $X^\tau$  denote the process  $X$  stopped at  $\tau$ , and we define the stochastic interval

$$\llbracket 0, \tau \rrbracket = \{(\omega, t) \in \Omega \times \mathbb{R}_+ : 0 \leq t < \tau(\omega)\}.$$

Note that stochastic intervals are disjoint from  $\Omega \times \{\infty\}$  by definition.

A process  $X$  on a stochastic interval  $\llbracket 0, \tau \rrbracket$ , where  $\tau$  is a stopping time, is the restriction to  $\llbracket 0, \tau \rrbracket$  of some process. In this paper,  $\tau$  will be a foretellable time; that is, a  $[0, \infty]$ -valued stopping time that admits a nondecreasing sequence  $(\tau_n)_{n \in \mathbb{N}}$  of stopping times, with  $\tau_n < \tau$  almost surely for all  $n \in \mathbb{N}$  on the event  $\{\tau > 0\}$ , and  $\lim_{n \uparrow \infty} \tau_n = \tau$  almost surely. Such a sequence is called an announcing sequence. Every predictable time is foretellable, and if the stochastic basis is complete the converse also holds; see Jacod and Shiryaev (2003, Theorem I.2.15 and I.2.16).

If  $\tau$  is a foretellable time and  $X$  is a process on  $\llbracket 0, \tau \rrbracket$ , we say that  $X$  is a semimartingale on  $\llbracket 0, \tau \rrbracket$  if there exists an announcing sequence  $(\tau_n)_{n \in \mathbb{N}}$  for  $\tau$  such that  $X^{\tau_n}$  is a semimartingale for each  $n \in \mathbb{N}$ . Local martingales and local supermartingales on  $\llbracket 0, \tau \rrbracket$  are defined analogously. Basic notions for semimartingales carry over by localization to semimartingales on stochastic intervals. For instance, if  $X$  is a semimartingale on  $\llbracket 0, \tau \rrbracket$ , its quadratic variation process  $[X, X]$  is defined as the process on  $\llbracket 0, \tau \rrbracket$  that satisfies  $[X, X]^{\tau_n} = [X^{\tau_n}, X^{\tau_n}]$  for each  $n \in \mathbb{N}$ . Its jump measure  $\mu^X$  is defined analogously, as are

stochastic integrals with respect to  $X$  (or  $\mu^X$ ). In particular,  $H$  is called  $X$ -integrable if it is  $X^{\tau_n}$ -integrable for each  $n \in \mathbb{N}$ , and  $H \cdot X$  is defined as the semimartingale on  $\llbracket 0, \tau \rrbracket$  that satisfies  $(H \cdot X)^{\tau_n} = H \cdot X^{\tau_n}$  for each  $n \in \mathbb{N}$ . We refer to Maisonneuve (1977), Jacod (1979), and Appendix A in Carr et al. (2014) for further details on local martingales on stochastic intervals.

The following version of the supermartingale convergence theorem is a useful technical tool for studying stochastic exponentials. It has been proven, for example, in Carr et al. (2014) via Doob's inequalities. For sake of completeness, we provide an alternative proof here.

**Lemma 2.1.** *If  $X$  is a local supermartingale on  $\llbracket 0, \tau \rrbracket$  bounded from below, then  $\lim_{t \uparrow \tau} X_t$  exists in  $\mathbb{R}$  and  $[X, X]_\tau = \lim_{t \uparrow \tau} [X, X]_t$  is finite.*

*Proof.* If  $\tau = \infty$  the statement is classical; see for instance (Dellacherie and Meyer, 1982, Theorem VI.6). The assertion regarding the quadratic variation follows from the fact that  $X$  can be extended to a supermartingale on  $[0, \infty]$ . Suppose now that  $\tau$  is a general foretellable time with announcing sequence  $(\tau_n)_{n \in \mathbb{N}}$ . Consider the time-change  $C = (C_u)_{u \geq 0}$  obtained as the right-continuous inverse of the strictly increasing càdlàg adapted process  $A$  on  $\llbracket 0, \tau \rrbracket$  given by

$$A_t = n + f(t - \tau_n), \quad \tau_n \leq t < \tau_{n+1}, \quad n \in \mathbb{N} \cup \{0\},$$

where  $f(t) = 1 - e^{-t}$  and  $\tau_0 = 0$ . The time-changed process  $Y = (X_{C_u})_{u \geq 0}$  is then a supermartingale with respect to the time-changed filtration, still of course bounded from below. Indeed, consider arbitrary  $u < v$ , and let  $n \in \mathbb{N}$  be such that  $v \in (n - 1, n]$ . Then  $C_u \leq C_v \leq \tau_n$ , and hence

$$\mathbb{E}[X_{C_v} \mid \mathcal{F}_{C_u}] = \mathbb{E}[X_{C_v}^{\tau_n} \mid \mathcal{F}_{C_u}] \leq X_{C_u}^{\tau_n} = X_{C_u}$$

by the optional sampling theorem. Thus  $Y$  is a supermartingale bounded from below, and therefore converges. Let now  $(t_k)_{k \in \mathbb{N}}$  be any sequence with  $t_k \uparrow \tau$ . Since  $A$  is strictly increasing, there is a sequence  $(u_k)_{k \in \mathbb{N}}$  with  $t_k = C_{u_k}$  and  $u_k \uparrow \infty$ . Thus  $\lim_{k \uparrow \infty} X_{t_k} = \lim_{k \uparrow \infty} Y_{u_k} = \lim_{u \uparrow \infty} Y_u$ , proving that the limit  $\lim_{t \uparrow \tau} X_t$  exists in  $\mathbb{R}$ . The statement regarding the quadratic variation follows from the fact that  $[X, X]_\tau = [Y, Y]_\infty < \infty$ .  $\square$

### 3 Stochastic exponentials and logarithms

In this section, we define stochastic exponentials and logarithms, develop some of their properties, and show that they are inverse to each other.

**Definition 3.1** (Stochastic exponential). Let  $\tau$  be a foretellable time, and let  $X$  be a semimartingale on  $\llbracket 0, \tau \rrbracket$  with  $\Delta X \geq -1$ . The *stochastic exponential* of  $X$  is the process

$\mathcal{E}(X)$  defined by

$$\mathcal{E}(X)_t = \exp\left(X_t - \frac{1}{2}[X^c, X^c]_t\right) \prod_{0 < s \leq t} (1 + \Delta X_s) e^{-\Delta X_s} \quad (3.1)$$

for all  $t < \tau$ , and by  $\mathcal{E}(X)_t = 0$  for all  $t \geq \tau$ .  $\square$

**Remark 3.2.** Note that the infinite product converges along every path where  $\Delta X \geq -1$ , since each factor lies in  $[0, 1]$ . Strictly speaking, the assumption  $\Delta X \geq -1$  (almost surely!) does not exclude larger negative jumps occurring on a nullset. A more pedantic definition would thus replace  $\Delta X$  by  $\Delta X \vee (-1)$  in (3.1). Furthermore, the condition  $\Delta X \geq -1$  could be dropped altogether, in which case the resulting stochastic exponential may change sign. This case is of limited use in applications, and we do not consider it here.  $\square$

The process  $\mathcal{E}(X)$  is sometimes also called generalized stochastic exponential; see, for example, Mijatović et al. (2012). If  $(\tau_n)_{n \in \mathbb{N}}$  is an announcing sequence for  $\tau$ , then  $\mathcal{E}(X)$  of Definition 3.1 coincides on  $\llbracket 0, \tau_n \rrbracket$  with the usual (Doléans-Dade) stochastic exponential of  $X^{\tau_n}$ . This shows that  $\mathcal{E}(X)$  coincides with the classical notion when  $\tau = \infty$ . Many properties of stochastic exponentials thus remain valid. For instance, if  $\Delta X > -1$  then  $\mathcal{E}(X)$  is strictly positive on  $\llbracket 0, \tau \rrbracket$ . If  $\Delta X_t = -1$  for some  $t \in [0, \tau)$  then  $\mathcal{E}(X)$  jumps to zero at time  $t$  and stays there. Also, on  $\llbracket 0, \tau \rrbracket$ ,  $\mathcal{E}(X)$  is the unique solution to the equation

$$Z = e^{X_0} + Z_- \cdot X \quad \text{on } \llbracket 0, \tau \rrbracket; \quad (3.2)$$

see Doléans-Dade (1976). We also record the more succinct expression

$$\mathcal{E}(X) = \mathbf{1}_{\llbracket 0, \tau \rrbracket} \exp\left(X - \frac{1}{2}[X^c, X^c] - (x - \log(1 + x)) * \mu^X\right), \quad (3.3)$$

where we use the convention  $-\log(0) = \infty$  and  $e^{-\infty} = 0$ .

We are particularly interested in the case where  $X$  is a local supermartingale on  $\llbracket 0, \tau \rrbracket$ . Then  $\mathcal{E}(X)$  is also a local supermartingale on  $\llbracket 0, \tau \rrbracket$  due to (3.2). Moreover, an application of the conditional Fatou lemma shows that  $\mathcal{E}(X)$  is in fact a supermartingale globally, i.e. on  $\llbracket 0, \infty \rrbracket$ . Indeed, for all  $0 \leq s < t$  we get

$$\begin{aligned} \mathbb{E}[\mathcal{E}(X)_t \mid \mathcal{F}_s] &= \mathbb{E}[\mathcal{E}(X)_t \mathbf{1}_{\{\tau > t\}} \mid \mathcal{F}_s] \\ &\leq \liminf_{n \rightarrow \infty} \mathbb{E}[\mathcal{E}(X)_{t \wedge \tau_n} \mathbf{1}_{\{\tau > t\}} \mid \mathcal{F}_s] \\ &\leq \liminf_{n \rightarrow \infty} \mathbb{E}[\mathcal{E}(X)_{s \wedge \tau_n} \mathbf{1}_{\{\tau > s\}}] \\ &= \mathcal{E}(X)_s. \end{aligned} \quad (3.4)$$

The following results characterizes the convergence to zero of  $\mathcal{E}(X)$  in terms of the behavior of  $X$ .

**Lemma 3.3.** *If  $X$  is a local supermartingale on  $\llbracket 0, \tau \rrbracket$  with  $\Delta X \geq -1$ , then the following set identity holds almost surely:*

$$\left\{ \lim_{t \uparrow \tau} \mathcal{E}(X)_t = 0 \right\} = \left\{ \lim_{t \uparrow \tau} X_t \text{ does not exist in } \mathbb{R} \right\} \cup \left\{ \Delta X_t = -1 \text{ for some } t \in [0, \tau) \right\}.$$

*Proof.* We argue the equality by showing the two inclusions.

“ $\supset$ ”: For an arbitrary  $m \in \mathbb{N}$ , define the stopping time  $\rho = \inf\{t \geq 0: X_t \leq -m\}$ . Then  $X^\rho$  is a local supermartingale on  $\llbracket 0, \tau \rrbracket$  bounded from below by  $-m - 1$ , whence  $\lim_{t \uparrow \tau} X_t^\rho$  exists in  $\mathbb{R}$  by Lemma 2.1. Since  $X$  coincides with  $X^\rho$  on  $\{X \geq -m\}$ , we deduce that

$$\left\{ \lim_{t \uparrow \tau} X_t \text{ does not exist in } \mathbb{R} \right\} \subset \bigcap_{m \in \mathbb{N}} \{X \geq -m\}^c \subset \left\{ \liminf_{t \uparrow \tau} X_t = -\infty \right\}.$$

Since (3.3) yields  $0 \leq \mathcal{E}(X) \leq \mathbf{1}_{\llbracket 0, \tau \rrbracket} e^X$  and the limit  $\lim_{t \uparrow \tau} \mathcal{E}(X)_t$  exists by Lemma 2.1, the inclusion follows.

“ $\subset$ ”: By taking complements, it suffices to pick  $m \in \mathbb{N}$  arbitrarily and argue that

$$\{X \geq -m\} \cap \{\Delta X > -1\} \subset \left\{ \inf_{t < \tau} X_t - \frac{1}{2} [X^c, X^c]_\tau - (x - \log(1+x)) * \mu_\tau^X > -\infty \right\}.$$

On the event on the left-hand side we have  $X = X^\rho$  with  $\rho$  as above, and thus  $[X, X]_\tau < \infty$  by Lemma 2.1. As a consequence,

$$(x - \log(1+x)) \mathbf{1}_{x > -1/2} * \mu_\tau^X \leq x^2 * \mu_\tau^X \leq [X, X]_\tau < \infty.$$

Finally, the fact that  $[X, X]_\tau < \infty$  also yields that  $\Delta X_t < -1/2$  occurs for at most finitely many  $t < \tau$ , whence  $\mathbf{1}_{x < -1/2} (x - \log(1+x)) * \mu_\tau^X < \infty$  on the event  $\{\Delta X > -1\}$ . This concludes the proof.  $\square$

We now consider the stochastic logarithm of a nonnegative semimartingale  $Z$  that stays at zero after reaching it. In preparation for this, recall that for a stopping time  $\rho$  and an event  $A \in \mathcal{F}$ , the *restriction of  $\rho$  to  $A$*  is given by

$$\rho(A) = \rho \mathbf{1}_A + \infty \mathbf{1}_{A^c}.$$

Here  $\rho(A)$  is a stopping time if and only if  $A \in \mathcal{F}_\rho$ . Define the following stopping times associated to a nonnegative semimartingale  $Z$  (recall our convention that  $Z_{0-} = Z_0$ ):

$$\begin{aligned} \tau_0 &= \inf\{t \geq 0 : Z_t = 0\}; \\ \tau_c &= \tau_0(A_C), \quad A_C = \{Z_{\tau_0-} = 0\}; \\ \tau_J &= \tau_0(A_J), \quad A_J = \{Z_{\tau_0-} > 0\}. \end{aligned} \tag{3.5}$$

These stopping times correspond to the two ways in which  $Z$  can reach zero: either continuously or by a jump. We have the following property of  $\tau_c$ .

**Lemma 3.4.** *Fix some nonnegative semimartingale  $Z$ . The stopping time  $\tau_c$  of (3.5) is foretellable.*

*Proof.* We claim that an announcing sequence  $(\sigma_n)_{n \in \mathbb{N}}$  for  $\tau_c$  is given by

$$\sigma_n = n \wedge \sigma'_n(A_n), \quad \sigma'_n = n \wedge \inf \left\{ t \geq 0 : Z_t \leq \frac{1}{n} \right\}, \quad A_n = \{Z_{\sigma'_n} > 0\}.$$

To prove this, we first observe that  $\sigma_n = n < \infty = \tau_c$  on  $A_n^c$  for all  $n \in \mathbb{N}$ . Moreover, we have  $\sigma_n = \sigma'_n < \tau_c$  on  $A_n$  for all  $n \in \mathbb{N}$ , where we used that  $Z_{\tau_c} = 0$  on the event  $\{\tau_c < \infty\}$ . We need to show that  $\lim_{n \uparrow \infty} \sigma_n = \tau_c$ . On the event  $A_C$ , see (3.5), we have  $\tau_c = \tau_0 = \lim_{n \uparrow \infty} \sigma'_n = \lim_{n \uparrow \infty} \sigma_n$  since  $A_C \subset A_n$  for all  $n \in \mathbb{N}$ . On the event  $A_C^c = \bigcup_{n=1}^{\infty} A_n^c$ , we have  $\tau_c = \infty = \lim_{n \uparrow \infty} n = \lim_{n \uparrow \infty} \sigma_n$ . Hence  $(\sigma_n)_{n \in \mathbb{N}}$  is an announcing sequence of  $\tau_c$ , as claimed.  $\square$

If a nonnegative semimartingale  $Z$  reaches zero continuously, the process  $H = \frac{1}{Z_-} \mathbf{1}_{\{Z_- > 0\}}$  explodes in finite time, and is therefore not left-continuous. In fact, it is not  $Z$ -integrable. However, if we view  $Z$  as a semimartingale on the stochastic interval  $\llbracket 0, \tau_c \rrbracket$ , then  $H$  is  $Z$ -integrable in the sense of stochastic integration on stochastic intervals. Thus  $H \cdot Z$  exists as a semimartingale on  $\llbracket 0, \tau_c \rrbracket$ , which we call the stochastic logarithm of  $Z$ .

**Definition 3.5** (Stochastic logarithm). Let  $Z$  be a nonnegative semimartingale with  $Z = Z^{\tau_0}$ . The semimartingale  $\mathcal{L}(Z)$  on  $\llbracket 0, \tau_c \rrbracket$  defined by

$$\mathcal{L}(Z) = \frac{1}{Z_-} \mathbf{1}_{\{Z_- > 0\}} \cdot Z \quad \text{on } \llbracket 0, \tau_c \rrbracket$$

is called the *stochastic logarithm* of  $Z$ .  $\square$

Our next goal is to clarify the relationship of stochastic exponentials and logarithms in the local supermartingale case. To this end, we make the following definition.

**Definition 3.6** (Maximality). Let  $\tau$  be a foretellable time, and let  $X$  be a semimartingale on  $\llbracket 0, \tau \rrbracket$ . We say that  $\tau$  is  $X$ -*maximal* if the inclusion

$$\{\tau < \infty\} \subset \left\{ \lim_{t \uparrow \tau} X_t \text{ does not exist in } \mathbb{R} \right\}$$

holds almost surely.  $\square$

Let now  $\mathfrak{J}$  be the set of all nonnegative supermartingales  $Z$  with  $Z_0 = 1$ . Any such process  $Z$  automatically satisfies  $Z = Z^{\tau_0}$ . Furthermore, let  $\mathfrak{L}$  denote the set of all stochastic processes  $X$  satisfying the following conditions:

- (i)  $X$  is a local supermartingale on  $\llbracket 0, \tau \rrbracket$  for some foretellable,  $X$ -maximal time  $\tau$ .
- (ii)  $X_0 = 0$ ,  $\Delta X \geq -1$  on  $\llbracket 0, \tau \rrbracket$ , and  $X$  is constant after the first time  $\Delta X = -1$ .

The next theorem extends the classical correspondence between strictly positive local martingales and local martingales with jumps strictly greater than  $-1$ . The reader is referred to Proposition I.5 in Lépingle and Mémin (1978) and Appendix A of Kardaras (2008) for related results.

**Theorem 3.7** (Relationship of stochastic exponential and logarithm). *The stochastic exponential  $\mathcal{E}$  is a bijection from  $\mathfrak{L}$  to  $\mathfrak{Z}$ , and its inverse is the stochastic logarithm  $\mathcal{L}$ . Suppose  $Z = \mathcal{E}(X)$  for some  $Z \in \mathfrak{Z}$  and  $X \in \mathfrak{L}$ . Then  $\tau = \tau_c$ , where  $\tau$  is the foretellable  $X$ -maximal time corresponding to  $X$ , and  $\tau_c$  is given by (3.5).*

*Proof.* First,  $\mathcal{E}$  maps each  $X \in \mathfrak{L}$  to some  $Z \in \mathfrak{Z}$  with  $\tau_c = \tau$ . Since  $Z = \mathcal{E}(X)$  is a nonnegative supermartingale by (3.4) with  $Z_0 = 1$ , we have  $Z \in \mathfrak{Z}$  and only need to argue that  $\tau_c = \tau$ . To see this, observe that  $\{\tau_c < \tau\}$  is a nullset due to (3.3) and the fact that  $X - \frac{1}{2}[X^c, X^c] - (x - \log(1 + x)) * \mu^X$  is a finite-valued càdlàg process on  $\llbracket 0, \tau \rrbracket$ . Thus  $\tau \leq \tau_c$ . Moreover, on  $\{\tau_0 < \tau < \tau_c\}$  we have  $\Delta X_{\tau_0} = -1$ , hence  $X$  is constant on  $\llbracket \tau_0, \tau \rrbracket$ , contradicting the  $X$ -maximality of  $\tau$ . Therefore, on  $\{\tau < \tau_c\}$  we have  $\lim_{t \uparrow \tau} \mathcal{E}(X)_t > 0$ . But this is incompatible with Lemma 3.3 and the  $X$ -maximality of  $\tau$ , unless  $\{\tau < \tau_c\}$  is a nullset. We conclude that  $\tau_c = \tau$ , as claimed.

Next,  $Z = \mathcal{E}(\mathcal{L}(Z))$  holds for each  $Z \in \mathfrak{Z}$ . Indeed,  $Z$  satisfies the equation  $Z = 1 + Z_- (Z_-)^{-1} \mathbf{1}_{\{Z_- > 0\}} \cdot Z = 1 + Z_- \cdot \mathcal{L}(Z)$  on  $\llbracket 0, \tau_c \rrbracket$ , whose unique solution is  $\mathcal{E}(\mathcal{L}(Z))$  on  $\llbracket 0, \tau_c \rrbracket$ .

Further,  $\mathcal{L}$  maps each  $Z \in \mathfrak{Z}$  to some  $X \in \mathfrak{L}$  with  $\tau = \tau_c$ . Indeed,  $X = \mathcal{L}(Z)$  is a local supermartingale on  $\llbracket 0, \tau_c \rrbracket$  with  $X_0 = 0$ ,  $\Delta X = Z/Z_- - 1 \geq -1$  on  $\llbracket 0, \tau_c \rrbracket$ , and  $X$  is constant after the first time  $\Delta X = -1$ . It remains to check that  $\tau = \tau_c$  is  $X$ -maximal. To this end, observe that  $Z = \mathcal{E}(\mathcal{L}(Z)) = \mathcal{E}(X)$ . Thus on  $\{\tau < \infty\}$  we have  $\tau_J = \infty$  and hence  $\Delta X > -1$ . It follows that

$$\{\tau < \infty\} \subset \{\lim_{t \uparrow \tau} Z_t = 0\} \cap \{\Delta X > -1\} \subset \{\lim_{t \uparrow \tau} X_t \text{ does not exist in } \mathbb{R}\},$$

where Lemma 3.3 yields the second inclusion. This shows that  $\tau$  is  $X$ -maximal as claimed.

So far we have proved that  $\mathcal{E}$  maps  $\mathfrak{L}$  into  $\mathfrak{Z}$ , that  $\mathcal{L}$  maps  $\mathfrak{Z}$  into  $\mathfrak{L}$ , and that  $\mathcal{E} \circ \mathcal{L} = \text{id}$ . Thus it remains to argue that  $\mathcal{L} \circ \mathcal{E} = \text{id}$ . To see this, let  $X \in \mathfrak{L}$  and use the definition of the stochastic logarithm along with (3.2) and the fact that  $\tau = \tau_c$  to obtain

$$\mathcal{L}(\mathcal{E}(X)) = \frac{1}{\mathcal{E}(X)_-} \mathbf{1}_{\{\mathcal{E}(X)_- > 0\}} \cdot \mathcal{E}(X) = \mathbf{1}_{\{\mathcal{E}(X)_- > 0\}} \cdot X = X \quad \text{on } \llbracket 0, \tau \rrbracket,$$

where the last equality follows since  $X$  is constant after the first time  $\Delta X = -1$ .  $\square$

## 4 Reciprocals of stochastic exponentials

Reciprocals of stochastic exponentials appear naturally in connection with changes of probability measure. We now develop some identities related to such reciprocals. The following

function plays an important role:

$$\phi : (-1, \infty) \rightarrow (-1, \infty), \quad \phi(x) = -1 + \frac{1}{1+x}.$$

Note that  $\phi$  is an involution, that is,  $\phi(\phi(x)) = x$ . The following notation is convenient: Given functions  $F : \Omega \times \mathbb{R}_+ \times \mathbb{R} \rightarrow \mathbb{R}$  and  $f : \mathbb{R} \rightarrow \mathbb{R}$ , we write  $F \circ f$  for the function  $(\omega, t, x) \mapsto F(\omega, t, f(x))$ . We now identify the reciprocal of a stochastic exponential or, more precisely, the stochastic logarithm of this reciprocal. Part of the following result is contained in Lemma 3.4 of Karatzas and Kardaras (2007).

**Theorem 4.1** (Reciprocal of a stochastic exponential). *Let  $\tau$  be a foretellable time, and let  $X$  be a semimartingale on  $\llbracket 0, \tau \rrbracket$  with  $\Delta X \geq -1$ . Define the semimartingale*

$$Y = -X + [X^c, X^c] + \frac{x^2}{1+x} \mathbf{1}_{x > -1} * \mu^X \quad \text{on } \llbracket 0, \tau \rrbracket. \quad (4.1)$$

*Then  $\mathcal{E}(X)\mathcal{E}(Y) = 1$  on  $\llbracket 0, \tau \wedge \tau_J \rrbracket$ . Furthermore, for any nonnegative function  $G : \Omega \times \mathbb{R}_+ \times \mathbb{R} \rightarrow \mathbb{R}_+$  we have*

$$G * \mu^Y = (G \circ \phi) * \mu^X \quad \text{on } \llbracket 0, \tau \wedge \tau_J \rrbracket. \quad (4.2)$$

**Remark 4.2.** Since  $|x^2/(1+x)| \leq 2x^2$  for  $|x| \leq 1/2$ , the process  $x^2/(1+x) * \mu^X$  appearing in (4.1) is finite-valued on  $\llbracket 0, \tau \rrbracket$ .  $\square$

**Remark 4.3.** Since  $\phi$  is an involution, the identity (4.2) is equivalent to

$$F * \mu^X = (F \circ \phi) * \mu^Y \quad \text{on } \llbracket 0, \tau \wedge \tau_J \rrbracket$$

for the nonnegative function  $F = G \circ \phi$ .  $\square$

*Proof of Theorem 4.1.* Note that we have

$$\Delta Y = -\Delta X + \frac{(\Delta X)^2}{1 + \Delta X} = \phi(\Delta X) \quad \text{on } \llbracket 0, \tau \wedge \tau_J \rrbracket.$$

This implies (4.2). Now, applying (4.2) to the function  $G(y) = y - \log(1+y)$  yields

$$(y - \log(1+y)) * \mu^Y = \left( -1 + \frac{1}{1+x} + \log(1+x) \right) * \mu^X \quad \text{on } \llbracket 0, \tau \wedge \tau_J \rrbracket.$$

A direct calculation then gives  $\mathcal{E}(Y) = 1/\mathcal{E}(X)$  on  $\llbracket 0, \tau \wedge \tau_J \rrbracket$ . This completes the proof.  $\square$

## 5 Examples

**Example 5.1.** Let  $X \in \mathfrak{L}$  be Brownian motion. Then the stopping time  $\tau = \infty$  is  $X$ -maximal, and  $\mathcal{E}(X)_t = e^{B_t - t/2}$  for all  $t \geq 0$  is geometric Brownian motion. From (4.1)  $Y_t = t - B_t$  for all  $t \geq 0$  satisfies  $\mathcal{E}(X)\mathcal{E}(Y) = 1$ .  $\square$

**Example 5.2.** Let  $E$  be a standard exponentially distributed random variable and assume that  $\mathbb{F}$  is the smallest right-continuous filtration such that  $\mathbf{1}_{[0, E]}$  is adapted. Let now  $X$  be given by  $X_t = t - \mathbf{1}_{\{\rho \leq t\}}$  for all  $t \geq 0$ . Then  $X \in \mathfrak{L}$ ,  $X$  is a martingale,  $\tau = \infty$  is  $X$ -maximal, and  $\mathcal{E}(X)_t = e^t \mathbf{1}_{[0, E]}$  for all  $t \geq 0$ .  $\square$

**Example 5.3.** Let  $(\Theta_n)_{n \in \mathbb{N}}$  denote a sequence of independent random variables with  $\mathbb{P}(\Theta_n = 1) = \mathbb{P}(\Theta_n = -1) = 1/2$  and let  $X_t = \sum_{n=1}^{[t]} \Theta_n$  for all  $t \geq 0$  be a standard random walk, where  $[t]$  denotes the integer part of  $t$ . Assume that  $\mathbb{F}$  is the smallest right-continuous filtration such that  $X$  is adapted. Define the stopping time

$$\rho = \inf\{t \geq 0 : \Delta X_t = -1\}.$$

Then  $\mathcal{E}(X)_t = \mathcal{E}(X^\rho)_t = 2^{[t]} \mathbf{1}_{[0, \rho]}$  for all  $t \geq 0$  and  $X^\rho = \mathcal{L}(\mathcal{E}(X)) \in \mathfrak{L}$ .  $\square$

**Example 5.4.** Similarly to Example 5.3, let  $(\Theta_n)_{n \in \mathbb{N}}$  denote a sequence of independent random variables with  $\mathbb{P}(\Theta_n = 1/2) = \mathbb{P}(\Theta_n = -1/2) = 1/2$ . Let now  $X$  be a local martingale on  $[0, 1)$ , given as a random walk that jumps at the deterministic times  $(t_n)_{n \in \mathbb{N}}$  where  $t_n = 1 - n^{-1}$ . That is,  $X_t = \sum_{n: t_n \leq t} \Theta_n$  for all  $t \in [0, 1)$ . Again, assume that  $\mathbb{F}$  is the smallest right-continuous filtration such that  $X$  is adapted. Then the deterministic stopping time  $\tau = 1$  is  $X$ -maximal; hence  $X \in \mathfrak{L}$ . Moreover, the stochastic exponential  $Z = \mathcal{E}(M) \in \mathfrak{Z}$  is given by

$$Z_t = \exp\left(\sum_{n: t_n \leq t} \log(1 + \Theta_n)\right) = \prod_{n: t_n \leq t} (1 + \Theta_n).$$

Theorem 3.7 yields that  $\tau_c = 1$ , where  $\tau_c$  was defined in (3.5). Alternatively, the strong law of large numbers and the fact that  $\mathbb{E}[\log(1 + \Theta_1)] < 0$  imply  $\mathbb{P}(\lim_{t \uparrow 1} Z_t = 0) = 1$ . This illustrates that  $\tau_c$  in (3.5) can be finite, even if the local martingale  $Z$  has no continuous component.

Define next the semimartingale  $Y$  on  $[0, 1]$  by

$$Y_t = -X_t + \frac{x^2}{1+x} * \mu_t^X = \left(-x + \frac{x^2}{1+x}\right) * \mu_t^X = \frac{-x}{1+x} * \mu_t^X \quad \text{for all } t \in [0, 1).$$

Theorem 4.1 yield that  $\mathcal{E}(X)\mathcal{E}(Y) = 1$  on  $[0, 1]$ . Indeed, using (3.3), we get

$$\begin{aligned} \mathcal{E}(Y) &= \mathbf{1}_{[0, 1]} \exp\left(\log(1+y) * \mu^Y\right) = \mathbf{1}_{[0, 1]} \exp\left(\log\left(1 + \frac{-x}{1+x}\right) * \mu^X\right) \\ &= \mathbf{1}_{[0, 1]} \exp\left(-\log(1+x) * \mu^X\right), \end{aligned}$$

which confirms this claim.  $\square$

**Example 5.5.** Let  $B$  be Brownian motion starting in zero, define the stopping time

$$\rho = \inf\{t \geq 0 : B_t = -1\}$$

and the nonnegative martingale  $Z = 1 + B^\rho \in \mathfrak{F}$ . That is,  $Z$  is Brownian motion started in one and stopped as soon as it hits zero. We now compute

$$X = \mathcal{L}(Z) = \frac{1}{Z} \mathbf{1}_{\{Z > 0\}} \cdot Z = \frac{1}{1+B} \cdot B \quad \text{on } \llbracket 0, \rho \rrbracket.$$

Note that  $\rho$  is indeed  $X$ -maximal by Theorem 3.7. □

## References

- Carr, P., Fisher, T., and Ruf, J. (2014). On the hedging of options on exploding exchange rates. *Finance Stoch.*, 18(1):115–144.
- Dellacherie, C. and Meyer, P.-A. (1982). *Probabilities and Potential. B*, volume 72 of *North-Holland Mathematics Studies*. North-Holland Publishing Co., Amsterdam. Theory of Martingales, Translated from the French by J. P. Wilson.
- Doléans-Dade, C. (1976). On the existence and unicity of solutions of stochastic integral equations. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 36(2):93–101.
- Jacod, J. (1979). *Calcul Stochastique et Problemes de Martingales*. Springer, Berlin.
- Jacod, J. and Shiryaev, A. N. (2003). *Limit Theorems for Stochastic Processes*. Springer, Berlin, 2nd edition.
- Kallsen, J. and Shiryaev, A. N. (2002). The cumulant process and Esscher’s change of measure. *Finance and Stochastics*, 6(4):397–428.
- Karatzas, I. and Kardaras, C. (2007). The numéraire portfolio in semimartingale financial models. *Finance and Stochastics*, 11(4):447–493.
- Kardaras, C. (2008). Balance, growth and diversity of financial markets. *Annals of Finance*, 4(3):369–397.
- Larsson, M. and Ruf, J. (2014). Convergence of local supermartingales and Novikov-Kazamaki-type conditions for processes with jumps. Preprint, arXiv:1411.6229.
- Lépingle, D. and Mémmin, J. (1978). Sur l’intégrabilité uniforme des martingales exponentielles. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 42:175–203.

- Maisonneuve, B. (1977). Une mise au point sur les martingales locales continues définies sur un intervalle stochastique. In *Séminaire de Probabilités, XI*, pages 435–445.
- Mijatović, A., Novak, N., and Urusov, M. (2012). Martingale property of generalized stochastic exponentials. In *Séminaire de Probabilités XLIV*, volume 2046 of *Lecture Notes in Math.*, pages 41–59. Springer, Heidelberg.
- Rheinländer, T. (2010). Stochastic exponentials. In Cont, R., editor, *Encyclopedia for Quantitative Finance*. Wiley.